

356 Sedge Warbler




Sedge Warbler. Spring (24-III).

SEDGE WARBLER (*Acrocephalus schoenobaenus*)

IDENTIFICATION

11-13 cm. Brown reddish upperparts, streaked dark; rufous rump; distinct pale supercilium; crown streaked dark; yellow-white underparts.



Sedge Warbler. Pattern of head and crown.

SIMILAR SPECIES

Recalls a **Reed Warbler**, without pale supercilium. Similar to **Aquatic Warbler** with a pale band along crown. **Moustached Warbler** has black crown, broader pale supercilium and white throat.



Reed Warbler



Moustached Warbler



Aquatic Warbler

SEXING

Plumage of both sexes alike.


AGEING

2 types of age can be recognized:

Juvenile with fresh feathers (check flight feathers and tail); iris dark brown; breast with dark brown spots.

Adult with worn plumage; iris pale brown or reddish; breast unspotted.

In **spring**, after the **prebreeding** moult, ageing is not possible using plumage characters.



Sedge Warbler. Ageing. Pattern of breast: left adult; right juvenile.

356 Sedge Warbler



S e d g e
W a r b l e r .
A g e i n g .
C o l o u r o f i r i s :
t o p a d u l t ;
b o t t o m
j u v e n i l e .



Sedge Warbler. Autumn. Juvenile (13-VIII).

MOULT



Partial **postbreeding** moult, including body feathers and the central pair of tail feathers, but no flight feathers. No **postjuvenile** moult. Both types of age have a complete **prebreeding** moult in wintering quarters.

PHENOLOGY

I	II	III	IV	V	VI	VII	VIII	IX	X	XI	XII

STATUS IN ARAGON

On passage, very common in marshlands of the Ebro Valley.



Sedge Warbler.
Crown pattern: top left in spring (24-III); top right adult in autumn (25-VIII); left juvenile (13-VIII).



Sedge Warbler. Autumn. Adult (25-VIII).



Sedge Warbler. Breast pattern: top left in spring (24-III); top right adult in autumn (25-VIII); left juvenile (13-VIII).

356 Sedge Warbler



S e d g e Warbler. Head pattern and iris colour: top in spring (30-III) middle adult in autumn (25-VIII) ; bottom juvenile (13-VIII).



Sedge Warbler. Tail pattern: top left in spring (24-III); top right adult in autumn (25-VIII); left juvenile (13-VIII).



Sedge Warbler. Spring: pattern of wing (24-III).



Sedge Warbler. Plumage wear: top left in spring (24-III); top right adult in autumn (25-VIII); left juvenile (13-VIII).



Sedge Warbler. Autumn. Adult: pattern of wing (25-VIII).

356 Sedge Warbler



Sedge Warbler. Autumn. Juvenile: pattern of wing (13-VIII).